

Materials List for:

Measurements of Motor Function and Other Clinical Outcome Parameters in Ambulant Children with Duchenne Muscular Dystrophy

Sara Nagy^{1,2}, Simone Schmidt^{1,2}, Patricia Hafner^{1,3}, Andrea Klein^{1,4,5}, Daniela Rubino-Nacht¹, Vanya Gocheva¹, Oliver Bieri⁶, Carole Vuillerot⁷, Ulrike Bonati^{1,2}, Dirk Fischer^{1,2,3}

¹Division of Pediatric Neurology, University Children's Hospital Basel (UKBB), University of Basel

²Department of Neurology, University of Basel Hospital

³Division of Neurology, Medical University Clinic, Kantonsspital Baselland

⁴Division of Pediatric Neurology, CHUV

⁵Division of Neuropaediatrics, Inselspital, University Children's Hospital Bern

⁶Department of Radiology, Division of Radiological Physics, University of Basel Hospital

⁷Hospices Civils de Lyon, Hôpital Femme-Mère-Enfant, L'Escale, Service de Médecine Physique et de Réadaptation Pédiatrique

Correspondence to: Sara Nagy at sara.nagy@ukbb.ch

URL: <https://www.jove.com/video/58784>

DOI: [doi:10.3791/58784](https://doi.org/10.3791/58784)

Materials

Name	Company	Catalog Number	Comments
Physiotherapy mat	-	-	should not be slippery; alternatively use a wide examination table
Cushions	-	-	-
Table	-	-	with adjustable height; it should allow the patient to rest forearms while seated and elbows flexed at 90°
Chair	-	-	with adjustable height if possible; it should allow the patient to touch the floor with the feet while seated with the hips and knees flexed at 90°
Stopwatch	-	-	-
CD or CD-ROM glued onto a piece of cardboard	-	-	-
10 coins	-	-	dimensions: 20 mm wide and 2 mm thick (10 euro cents or equivalent)
Lead pencil	-	-	-
Tennis ball	-	-	-
Sheets of A4 paper or equivalent	-	-	weight: 70-80g
Clipboard	-	-	-
Two small traffic cones	-	-	-
Tape	-	-	for marking arrows and stop
Line traced on the floor	-	-	2 centimeters wide and 6 meters long
Corridor	-	-	indoor, straight, up to 30 meters long