

Materials List for:

Performing and Processing FNA of Anterior Fat Pad for Amyloid

Vinod B. Shidham^{1,2}, Bryan Hunt¹, Safwan S. Jaradeh³, Alexandru C. Barboi³, Sumana Devata⁴, Parameswaran Hari⁵

¹Department of Pathology, Medical College of Wisconsin

²Current Address: Department of Pathology, Wayne State University School of Medicine Detroit Medical Center

³Department of Neurology, Medical College of Wisconsin

⁴Department of Medicine, Medical College of Wisconsin

⁵Division of Neoplastic Diseases and Related Disorders, Medical College of Wisconsin

Correspondence to: Vinod B. Shidham at vshidham@med.wayne.edu

URL: <https://www.jove.com/video/1747>

DOI: [doi:10.3791/1747](https://doi.org/10.3791/1747)

Materials

Name	Company	Catalog Number	Comments
Alcohol swabs			
gauze pads			
marking pen			
10 mL syringes x2			
1% lidocaine			local anesthetic- If lidocaine is used alone, it causes an initial burning sensation after instillation. This can be prevented by using a 1:1 mixture of 1% lidocaine and 1% sodium-bi-carbonate
18 gauge(G) 1 ½ inch needles x2 (for performing FNAB)			
25G 1 ½ inch needle			for injecting local anesthetic
FNA syringe holder ("gun")			
bandage			
vial of glutaraldehyde			for electron microscopy
biopsy container of 10% formalin			