

**Materials List for:**

# Subcutaneous Angiotensin II Infusion using Osmotic Pumps Induces Aortic Aneurysms in Mice

Hong Lu<sup>1</sup>, Deborah A. Howatt<sup>1</sup>, Anju Balakrishnan<sup>1</sup>, Jessica J. Moorlegghen<sup>1</sup>, Debra L. Rateri<sup>1</sup>, Lisa A. Cassis<sup>2</sup>, Alan Daugherty<sup>1</sup>

<sup>1</sup>Saha Cardiovascular Research Center, University of Kentucky

<sup>2</sup>Department of Pharmacology and Nutritional Sciences, University of Kentucky

Correspondence to: Alan Daugherty at [adaugh@uky.edu](mailto:adaugh@uky.edu)

URL: <https://www.jove.com/video/53191>

DOI: [doi:10.3791/53191](https://doi.org/10.3791/53191)

## Materials

Name	Company	Catalog Number	Comments
Angiotensin II	Bachem	H-1705	compound used to induce aortic aneurysms
Alzet Osmotic Pumps	DURECT Corporation	Alzet Model 2004	feasible for 28-day infusion in mice weighed > 20 g
Saturated fat-enriched diet	Harlan Teklad	TD.88137	42% calories/calories to stimulate hypercholesterolemia in LDL receptor -/- mice